

to say, however, to what extent the colours may have been altered by the spirit: some of the scales also may have been rubbed off.

CHROMIS FACETUS. Jen.

*C. supra virescenti-niger, lateribus pallidioribus: dorso modice arcuato; fronte elevato, rostro summo ante oculos paululum excavato: limbo preoperculi poris quatuor conspicuis impresso: squamis latis, marginibus liberis levissime ciliatis: spinis dorsalibus quindecim, analibus sex: pinnis ventralibus longe acuminatis, ad analem pertingentibus: pinnâ caudali subæquali.*

D. 15/10; A. 6/3; C. 16, &c.; P. 14; V. 1/5.

LONG. unc. 5. lin. 9.

FORM.—Oblong-oval, very much compressed; the back moderately elevated, and more curved than the abdomen. Greatest depth a little behind the insertion of the pectorals, and contained twice and three-quarters in the entire length: thickness about two-fifths of the depth. Fore-head high: profile falling very obliquely, and slightly hollowed out in front of the eyes; the upper and under profile meeting at the mouth at nearly a right angle. Head contained not quite four times in the entire length; its own length and height nearly equal. Mouth small, protractile: jaws about equal, the lower one, if anything, a little the longest: lips not very fleshy. Maxillary rather slender, retiring almost entirely, when the mouth is closed, beneath the sub-orbital, the anterior margin of which is slightly hollowed out, and somewhat sinuous. Teeth in card in both jaws, forming a narrow band; the outermost row longer and stronger than the others, especially the four or six middle ones in front, which are somewhat conical and slightly hooked. Pharyngean teeth present, but none on the vomer or palatines. Eyes rather small, their diameter about one-fifth the length of the head; high in the cheeks, and a little nearer to the snout than to the posterior margin of the opercle: the space between broad, equalling nearly two diameters and-a-half. Nostrils consisting of a single round orifice half-way between the eye and the end of the snout. Preopercle with the basal margin short, and forming a slightly obtuse angle with the ascending one, the margin of which is entire. Opercle of a triangular form, broad at top, but narrowing off towards the bottom. Subopercle and interopercle much developed; their outer margins, taken together, rounded off nearly in a semicircle. Branchial membrane quite free all round, unattached to the isthmus, and but slightly emarginate. Snout, suborbital, jaws, and limb of the preopercle, naked; but the cheeks and rest of the opercular pieces scaly: the scales on the subopercle large. Four large pores on the limb of the preopercle, preceded by three others beneath the lower jaw: similar pores beneath the eye, and extending partially round it; one on the crown of the head, and a few smaller ones scattered about the snout; a large one just above the opercle, and another higher up on each side of the nape.

Scales on the body large; about twenty-five or twenty-six in a longitudinal row, and eleven or twelve in the depth; broader than long, with the free edges very minutely ciliated, the concealed portions with a fan of thirteen striæ, and the basal margins with twelve distinct

crenatures. Lateral line interrupted; its first portion at the depth of two and a half rows of scales beneath the dorsal, and stopping beneath the commencement of the soft part of that fin; recommencing three rows lower down, exactly in the middle of the depth, whence it runs straight to the caudal.

Dorsal commencing above the opercle; the spinous portion of nearly uniform height, and scarcely more than one-fifth of the depth; the soft portion much higher, and terminating in a sharp point behind. Anal answering to the posterior half of the dorsal, terminating opposite to it, and similarly pointed; with six spines, which, as well as the dorsal spines, are furnished with very conspicuous filamentous tags. Caudal nearly even. Pectorals rounded, but not very broad, their length rather more than three-fourths that of the head; the rays rather slender. Ventrals pointed; the first soft ray elongated, and reaching to the anal when laid back; the last ray attached at its base by a membrane to the abdomen. Rows of small scales between the rays of the caudal at the base of the fin; and a few small ones along the base of the dorsal and anal, more particularly on the soft portions.

COLOUR.—“Above, greenish black; the sides paler; slightly iridescent.”—D.—In spirits it appears of a nearly uniform brown all over, fins included.

Habitat, Maldonado, Rio Plata.

Mr. Darwin obtained this species at Maldonado, in a lake of fresh water, said sometimes to be a little brackish. It appears to belong to the genus *Chromis* of Cuvier, placed by him amongst the *Labridæ*, but having evidently very strong affinities to some of the *Sciænidæ*. It differs essentially from the *C. Brasiliensis* of Quoy and Gaimard,\* in having six anal spines, and being destitute of all markings and spots. I am not aware that it is described by any author.

1. SCARUS CHLORODON. Jen.

PLATE XXI.

*S. æruginoso cyaneus, capite et pinnis flavo-vittatis: maxillis exteriùs lævibus, marginibus crenatis; caninis ad angulos oris nullis: fronte gradatim proclivi: lineâ laterali tubis parum ramosis: squamis ubique striato-granulatis: pectoralibus acuminatis, radiis superioribus arcuatis: caudali radiis externis cæteris longioribus, acuminatis.*

D. 9/10; A. 3/9; C. 13. &c.; P. 15 vel 16; V. 1/5.

LONG. unc. 16.

FORM.—Of an oval form; the greatest depth one-third of the length, caudal excluded: dorsal and ventral lines equally convex. Head a little less than the depth of the body, not gibbous in front, but with the profile falling regularly and gradually from the commencement of the dorsal. Snout rather pointed. Jaws equal, their outer surface smooth, but crenated on their cutting edges. No spinous canines at the corners of the mouth. Eyes rather small, their diameter not one-fifth the length of the head, situate above the middle of the cheek, but equidistant from the posterior lobe of the opercle and the extremity of the snout. Snout in front of the eyes, and

\* Freycinet Voyage, (Zoologie) p. 286.